



On Isolated Points of the Spectrum of a Bounded Linear Operator

Christoph Schmoege

Proceedings of the American Mathematical Society, Vol. 117, No. 3. (Mar., 1993), pp. 715-719.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9939%28199303%29117%3A3%3C715%3AOIPOTS%3E2.0.CO%3B2-8>

Proceedings of the American Mathematical Society is currently published by American Mathematical Society.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/ams.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is an independent not-for-profit organization dedicated to creating and preserving a digital archive of scholarly journals. For more information regarding JSTOR, please contact support@jstor.org.

<http://www.jstor.org/>
Mon May 15 09:44:04 2006

ON ISOLATED POINTS OF THE SPECTRUM OF A BOUNDED LINEAR OPERATOR

CHRISTOPH SCHMOEGER

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. For a bounded linear operator A on a Banach space we characterize the isolated points in the spectrum of A , the Riesz points of A , and the poles of the resolvent of A .

1. TERMINOLOGY AND INTRODUCTION

Throughout this paper E will be an infinite-dimensional complex Banach space and A will be a bounded linear operator on E . We denote by $N(A)$ the kernel and by $A(E)$ the range of A . The spectrum of A will be denoted by $\sigma(A)$. The resolvent set $\varrho(A)$ of A is the complement of $\sigma(A)$ in the complex plane \mathbb{C} . For any λ in $\varrho(A)$ the resolvent operator $(\lambda I - A)^{-1}$ is denoted by $R_\lambda(A)$.

Let λ_0 be an isolated point in $\sigma(A)$. The spectral projection corresponding to λ_0 will be denoted by P_{λ_0} . We have $E = P_{\lambda_0}(E) \oplus N(P_{\lambda_0})$.

In [3] Mbekhta introduced two important subspaces of E :

$$\begin{aligned} K(A) = \{x \in E : \text{there exist } c > 0 \text{ and a sequence } (x_n)_{n \geq 1} \subseteq E \\ \text{such that } Ax_1 = x, \ Axx_{n+1} = x_n \text{ for all } n \in \mathbb{N}, \\ \text{and } \|x_n\| \leq c^n \|x\| \text{ for all } n \in \mathbb{N}\}, \end{aligned}$$

$$H_0(A) = \left\{x \in E : \lim_{n \rightarrow \infty} \|A^n x\|^{1/n} = 0\right\}$$

and proved the following

Theorem 1. *A point $\lambda_0 \in \sigma(A)$ is isolated in $\sigma(A)$ if and only if there is a bounded projection P on E such that*

$$P(E) = H_0(\lambda_0 I - A) \quad \text{and} \quad N(P) = K(\lambda_0 I - A).$$

In the present paper we shall prove that $\lambda_0 \in \sigma(A)$ is an isolated point of $\sigma(A)$ if and only if $K(\lambda_0 I - A)$ is closed and $E = K(\lambda_0 I - A) \oplus H_0(\lambda_0 I - A)$ (where \oplus denotes the algebraically direct sum). This characterization leads to

Received by the editors March 4, 1991 and, in revised form, June 24, 1991.

1991 *Mathematics Subject Classification.* Primary 47A10; Secondary 47B06.

Key words and phrases. Local spectral theory, isolated points of the spectrum, Riesz points.

©1993 American Mathematical Society
0002-9939/93 \$1.00 + \$.25 per page

a characterization of the poles of the resolvent of A and to a characterization of the Riesz points of A . This will be done in §3 of this paper.

2. PRELIMINARY RESULTS

The operator A is said to have the *single-valued extension property* (SVEP) in $\lambda_0 \in \mathbb{C}$ if for any holomorphic function $f: U \rightarrow E$, where U is a neighbourhood of λ_0 , with $(\lambda I - A)f(\lambda) \equiv 0$, the result is $f(\lambda) \equiv 0$. We say that A has the SVEP if A has the SVEP in each $\lambda \in \mathbb{C}$.

The following theorem collects some results due to Mbekhta (see [4]).

Theorem 2. (a) $A(K(A)) = K(A)$ and $A(H_0(A)) \subseteq H_0(A)$;
 (b) A has the SVEP in λ_0 if $H_0(\lambda_0 I - A)$ is closed;
 (c) A has the SVEP in λ_0 if and only if $K(\lambda_0 I - A) \cap H_0(\lambda_0 I - A) = \{0\}$.

The proof of the next result is immediate.

Proposition 1. Let $x \in H_0(A)$ and define the function g on $\mathbb{C} \setminus \{0\}$ by

$$g(\lambda) = \sum_{n=0}^{\infty} \frac{A^n x}{\lambda^{n+1}}.$$

Then g is holomorphic and $(\lambda I - A)g(\lambda) = x$ for all $\lambda \in \mathbb{C} \setminus \{0\}$.

Proposition 2. Let F be a closed subspace of E such that $A(F) = F$. Then $F \subseteq K(A)$.

Proof. Since F is a Banach space and $A(F) = F$, the open mapping theorem shows the existence of a constant $c > 0$ so that

$$(2.1) \quad \text{for each } u \in F \text{ there exists } v \in F \text{ such that} \\ Av = u \quad \text{and} \quad \|v\| \leq c\|u\|.$$

Let $x \in F$. Use (2.1) to construct a sequence $(x_n)_{n \geq 1} \subseteq F$ such that $Ax_1 = x$, $Ax_{n+1} = x_n$, and $\|x_n\| \leq c^n \|x\|$. It follows that $x \in K(A)$. \square

Let us review the classical definitions of ascent and descent. The *ascent* $p(A)$ and the *descent* $q(A)$ are the extended integers given by

$$p(A) = \inf\{n \geq 0 : N(A^n) = N(A^{n+1})\}, \\ q(A) = \inf\{n \geq 0 : A^n(E) = A^{n+1}(E)\}.$$

The infimum over the empty set is taken to be ∞ . It follows from [2, Satz 72.3] that if $p(A)$ and $q(A)$ are both finite then they are equal.

We have the following characterization of the poles of the resolvent of A (see [2, Satz 101.2]):

Theorem 3. The complex number λ_0 is a pole of $R_\lambda(A)$ if and only if $0 < p(\lambda_0 I - A) = q(\lambda_0 I - A) < \infty$. In this case we have

$$P_{\lambda_0}(E) = N((\lambda_0 I - A)^p) \quad \text{and} \quad N(P_{\lambda_0}) = (\lambda_0 I - A)^p(E),$$

where $p = p(\lambda_0 I - A)$ is the order of the pole λ_0 .

The next proposition is a generalization of [1, Theorem 2].

Proposition 3. Suppose that A has the SVEP in $\lambda_0 = 0$ and $q(A) < \infty$. Then $p(A) = q(A)$.

Proof. Let $q = q(A)$, $B = A^q$, and $\widehat{E} = E/N(B)$. Since $N(B)$ is closed, \widehat{E} is a Banach space. Let $\widehat{B}: \widehat{E} \rightarrow E$ be the corresponding canonical injection. It is easy to see that the operator $\widehat{B}^{-1}: A^q(E) \rightarrow \widehat{E}$ is closed, thus $A^q(E)$ is the domain of a closed linear operator. Since $A(A^q(E)) = A^q(E)$ and A has the SVEP in 0, [1, Corollary 4] shows that $N(A) \cap A^q(E) = \{0\}$. Use [2, Satz 72.1] to derive $p(A) < \infty$. \square

Corollary 1. The following assertions are equivalent:

- (a) 0 is a pole of $R_\lambda(A)$;
- (b) A has the SVEP in 0 and $q(A) < \infty$.

Proof. (a) implies (b). Since 0 is isolated in $\sigma(A)$, A has the SVEP in 0. Theorem 3 shows that $q(A) < \infty$.

(b) implies (a). Proposition 3 and Theorem 3. \square

3. ISOLATED POINTS OF THE SPECTRUM

The starting point of our investigation is

Proposition 4. Suppose that 0 is an isolated point in $\sigma(A)$. Then

- (a) $P_0(E) = H_0(A)$;
- (b) $N(P_0) = K(A)$.

Proof. (a) follows from [2, Satz 100.2].

(b) Since 0 is isolated in $\sigma(A)$, $\sigma(A|_{P_0(E)}) = \{0\}$ and $0 \in \varrho(A|_{N(P_0)})$ [2, Satz 100.1]. Then $N(P_0)$ is closed and $A(N(P_0)) = N(P_0)$. Hence, by Proposition 2, $N(P_0) \subseteq K(A)$. By Theorem 2(c), $K(A) \cap H_0(A) = \{0\}$. Therefore,

$$\begin{aligned} K(A) &= K(A) \cap E = K(A) \cap [N(P_0) \oplus P_0(E)] \\ &= N(P_0) + K(A) \cap H_0(A) = N(P_0). \quad \square \end{aligned}$$

Theorem 4. The following assertions are equivalent:

- (a) 0 is an isolated point in $\sigma(A)$;
- (b) $K(A)$ is closed and $E = K(A) \oplus H_0(A)$ (\oplus denotes the algebraically direct sum).

Proof. (a) implies (b). Use Proposition 4 or Theorem 1.

(b) implies (a). Since $K(A)$ is closed, $A(K(A)) = K(A)$ (Theorem 2(a)), and $N(A) \subseteq H_0(A)$, the operator $A: K(A) \rightarrow K(A)$ is invertible. Hence there exists $\varepsilon > 0$ such that $\lambda I - A|_{K(A)}$ is invertible if $|\lambda| < \varepsilon$. In particular,

$$(3.1) \quad (\lambda I - A)(K(A)) = K(A) \quad \text{if } |\lambda| < \varepsilon.$$

Since for all $\lambda \neq 0$, $N(\lambda I - A) \subseteq K(A)$, we have

$$(3.2) \quad N(\lambda I - A) = \{0\} \quad \text{if } 0 < |\lambda| < \varepsilon.$$

By Proposition 1, for all $\lambda \neq 0$,

$$(3.3) \quad H_0(A) \subseteq (\lambda I - A)(E).$$

Now, (3.1) and (3.3) imply

$$E = K(A) \oplus H_0(A) \subseteq (\lambda I - A)(E) \quad \text{if } 0 < |\lambda| < \varepsilon.$$

Consequently, $\{\lambda \in \mathbb{C} : 0 < |\lambda| < \varepsilon\} \subseteq \varrho(A)$ and the proof is complete. \square

Now we are in a position to present the announced characterization of the poles of the resolvent of A .

Theorem 5. *The following assertions are equivalent:*

- (a) 0 is a pole of the resolvent of A ;
- (b) A has the SVEP in 0 and $q(A) < \infty$;
- (c) There exists $p \in \mathbb{N}$ such that

$$N(A^p) = H_0(A) \quad \text{and} \quad A^p(E) = K(A);$$

- (d) A has the SVEP in 0 and there exists $p \in \mathbb{N}$ such that $K(A) = A^p(E)$;
- (e) $q(A) < \infty$ and $H_0(A)$ is closed.

Proof. By Corollary 1, (a) and (b) are equivalent.

(a) implies (c). Use Theorem 3 and Proposition 4.

(c) implies (a). By Theorem 3, we have to show that $p(A)$ and $q(A)$ are both finite. Since

$$N(A^{p+1}) \subseteq H_0(A) = N(A^p) \subseteq N(A^{p+1}),$$

we have $p(A) \leq p$. Use Theorem 2(a) to derive $A^{p+1}(E) = A(A^p(E)) = A(K(A)) = K(A) = A^p(E)$. Thus $q(A) \leq p$.

(a) implies (d). Use (b) and (c).

(d) implies (b). As in the proof of “(c) implies (a),” we have $A^p(E) = A^{p+1}(E)$, hence $q(A) < \infty$.

(a) implies (e). Clear.

(e) implies (b). By Theorem 2(b), A has the SVEP in 0 . \square

The remainder of this paper deals with Riesz points and Riesz operators. A complex number λ_0 is called a *Riesz point* of A , if

$$p(\lambda_0 I - A) = q(\lambda_0 I - A) < \infty \quad \text{and} \quad \dim N(\lambda_0 I - A) = \operatorname{codim}(\lambda_0 I - A)(E) < \infty.$$

Note that a Riesz point of A is either a pole of the resolvent (and hence isolated in $\sigma(A)$) or a point in the resolvent set $\varrho(A)$.

Proposition 5. *The complex number $\lambda_0 \in \sigma(A)$ is a Riesz point of A if and only if λ_0 is isolated in $\sigma(A)$ and the corresponding spectral projection is finite dimensional.*

Proof. [2, Satz 105.3]. \square

The next theorem uses the subspaces $K(A)$ and $H_0(A)$ and the SVEP to characterize the Riesz points of A .

Theorem 6. *The following assertions are equivalent:*

- (a) 0 is a Riesz point of A ;
- (b) $K(A)$ is closed, $\dim H_0(A) < \infty$, and $E = K(A) \oplus H_0(A)$, where \oplus denotes the algebraically direct sum;
- (c) $q(A) < \infty$ and $\dim H_0(A) < \infty$;
- (d) $\dim H_0(A) < \infty$ and $K(A) = A^p(E)$ for some $p \in \mathbb{N}$;
- (e) A has the SVEP in 0 , $q(A) < \infty$, and $\dim N(A) < \infty$.

Proof. (a) \Leftrightarrow (b). Proposition 4 and Theorem 4.

(c) \Rightarrow (a). Since $N(A^n) \subseteq N(A^{n+1}) \subseteq H_0(A)$ and $\dim H_0(A) < \infty$, there exists $p \in \mathbb{N}$ such that $\dim N(A^p) = \dim N(A^{p+1}) < \infty$. This gives $N(A^p) = N(A^{p+1})$, thus $p(A) < \infty$. By Theorem 3, 0 is a pole of $R_\lambda(A)$, hence 0 is isolated in $\sigma(A)$. Proposition 4 shows that $\dim P_0(E) < \infty$. Now use Proposition 5.

(a) \Rightarrow (d). Propositions 4 and 5 show that $\dim P_0(E) = \dim H_0(A) < \infty$ and $N(P_0) = K(A)$. Since 0 is a pole of $R_\lambda(A)$, we conclude from Theorem 3 that $K(A) = A^p(E)$ for some $p \in \mathbb{N}$.

(d) \Rightarrow (c). $A^{p+1}(E) = A(K(A)) = K(A) = A^p(E)$, thus $q(A) < \infty$.

(a) \Rightarrow (e). Clear.

(e) \Rightarrow (a). By Proposition 3, $p(A) = q(A) < \infty$. [2, Satz 72.6] shows that $\dim N(A) = \text{codim } A(E) < \infty$, thus 0 is a Riesz point of A . \square

The operator A is called a *Riesz operator* if every $\lambda \in \sigma(A) \setminus \{0\}$ is a Riesz point of A .

An immediate consequence of Theorem 6 is

Theorem 7. *The following assertions are equivalent:*

- (a) A is a Riesz operator;
- (b) $\dim H_0(\lambda I - A) < \infty$, $E = K(\lambda I - A) \oplus H_0(\lambda I - A)$, and $K(\lambda I - A)$ is closed for all $\lambda \in \sigma(A) \setminus \{0\}$;
- (c) $q(\lambda I - A) < \infty$ and $\dim H_0(\lambda I - A) < \infty$ for all $\lambda \in \sigma(A) \setminus \{0\}$;
- (d) $\dim H_0(\lambda I - A) < \infty$ for all $\lambda \in \sigma(A) \setminus \{0\}$ and for each $\lambda \in \sigma(A) \setminus \{0\}$ there exists $p(\lambda) \in \mathbb{N}$ such that $K(\lambda I - A) = (\lambda I - A)^{p(\lambda)}(E)$.

ACKNOWLEDGMENT

The author wishes to thank the referee for his suggestions that led to a simplification of the proof of Theorem 4.

REFERENCES

1. J. K. Finch, *The single-valued extension property on a Banach space*, Pacific J. Math. **58** (1975), 61–69.
2. H. Heuser, *Funktionalanalysis*, 2nd ed., Teubner, Stuttgart, 1986.
3. M. Mbekhta, *Généralisation de la décomposition de Kato aux opérateurs paranormaux et spectraux*, Glasgow Math. J. **29** (1987), 159–175.
4. —, *Sur la théorie spectrale locale et limite des nilpotents*, Proc. Amer. Math. Soc. **110** (1990), 621–631.

MATHEMATISCHES INSTITUT 1, UNIVERSITÄT KARLSRUHE, ENGLERSTR. 2, D-7500 KARLSRUHE 1, GERMANY